
Changes in biochemical and antioxidant enzymes activities play significant role in drought tolerance in soybean

Mishra, N.¹, Tripathi, M. K.^{1*}, Tripathi, N.², Tiwari, S.¹, Gupta, N.¹, Sharma, A.¹ and Shrivastav, M. K.³

¹Department of Plant Molecular Biology and Biotechnology, College of Agriculture, Rajmata Vijayaraje Scindia Agricultural University, Gwalior 474002, India; ²Directorate of Research Services, Jawaharlal Nehru Agricultural University, Jabalpur 482004, India; ³Department of Plant Breeding & Genetics, Jawaharlal Nehru Agricultural University, Jabalpur 482004, India.

Mishra, N., Tripathi, M. K., Tripathi, N. Tiwari, S., Gupta, N., Sharma, A. and Shrivastav, M. K. (2021). Changes in biochemical and antioxidant enzymes activities play significant role in drought tolerance in soybean. *International Journal of Agricultural Technology* 17(4):1425-1446.

Abstract Soybean genotypes were grouped into two clusters on the basis of biochemical profiling, anti-oxidant enzyme activities and protein profiling. Biochemical and antioxidant enzyme activity analysis among 53 genotypes revealed the presence of drought tolerance traits in three genotypes *viz.*, JS97-52, RVS-14 and JS95-60. The result obtained may contribute towards improvement of soybean genotypes with the development of drought tolerant varieties with the applications of conventional as well as molecular breeding approaches. These findings also provided a base for further research to investigate the drought tolerance mechanism in soybean crop using advanced biotechnological tools.

Keywords: Antioxidant enzymes, Biochemical parameters, Drought, Protein profiling, Sustainable agriculture

Introduction

Changes in environmental conditions are responsible for evolution of new challenges. Among these, climate change is an important issue which considerably affects agriculture. To achieve the target of grain production there is a big demand of water for irrigation of crops in most of the parts of the world. Due to this drought will be a big challenge in coming days for survival of more than 40% of people belongs to 54 countries (Gardner-Outlaw and Engelman, 1997). A crop species or genotype that is pliant to low rainfall intensity and unpredictable distribution and elevated temperature would have a decisive significance for sustainable food supply to the enduringly increasing world population. To breed a cultivar for such suboptimal environment by

* **Corresponding Author:** Tripathi, M. K.; **Email:** drmanojtripathi64@gmail.com

means of cautiously crafted biotechnological strategies is expected to sustain food security in inhospitable climates.

Soybean is also acknowledged as a 'miracle crop' due to over 40% protein and 20% oil (Tripathi and Tiwari, 2004; Tiwari & Tripathi, 2005; Mishra *et al.*, 2020; Upadhyay *et al.*, 2020a; Upadhyay *et al.*, 2020b). It needs an adequate water supply for the duration of its growth and development course to accomplish better yields (Buezo *et al.*, 2018). The plants of soybean have been found to be affected by drought at every stage of life (Dhanda *et al.*, 2004; Kachare, 2017; Kachare *et al.*, 2019; Mishra *et al.*, 2021a; Sharma *et al.*, 2021). Significant reductions in the levels of chlorophyll a, b, and total chlorophyll have been observed due to drought in soybean crop (Wu and Zhang, 2019). Plants develop various mechanisms to fight different stresses (Specht *et al.*, 2001) and these mechanisms may be due to alteration in biochemical pathways. Numerous biochemical parameters have been exploited to recognize tolerant genotype (s) to drought (Wang *et al.*, 2019; Sahu *et al.*, 2020; Choudhary *et al.*, 2021).

To distinguish the desired genotype(s), various biochemical parameters that are being utilized for selection are proline, membrane stability, total sugar, MDA, protein, antioxidant activities (catalase, glutathione reductase and peroxidase) etc. Earlier, it is reported that the antioxidant enzymes play major role in control, and removal of reactive oxygen species (ROS). The increased activities of superoxide dismutase (SOD), peroxidase (POD) and catalase (CAT) have been observed in crops as well as soybean under water stress (Kachare, 2017; Kachare *et al.*, 2019; Wang *et al.*, 2019). Enhancement of MDA content has also been recorded under water deficit in crops (Wang *et al.*, 2019; Sahu *et al.*, 2020).

Identification of drought tolerant genotypes due to biochemical alterations may provide a basis for the development of new plant varieties using conventional (Mishra *et al.*, 2021b) as well as molecular breeding approaches to fight water stress. The current investigation was executed to monitor drought tolerant soybean genotypes based on manifestation of different biochemical parameters, antioxidant enzymes activities and protein profile.

Materials and methods

The present investigation was consisted of 53 soybean genotypes (Table 1) with diverse reactions to drought *viz.*, susceptible and tolerant. The seeds were obtained from College of Agriculture, JNKVV, Jabalpur, RAK, College, Sehore and Zonal Agricultural Research Station, Morena, Madhya Pradesh, India.

Table 1. List of soybean genotypes with their parentage

S. No.	Genotypes	Source/Pedigree	S. No.	Genotypes	Source/Pedigree
1.	JS 20-29	JS 97-52 x JS 95-56	28.	RSC-10-52	NRC 37X JS335
2.	JS 20-69	JS 97-52 x SL 710	29.	SL -1123	Selection from AGS751
3.	JS 335	JS 78-77 x JS 71-05	30.	SL-1068	SL755XSL525
4.	JS 20-98	JS 97-52x JS SL710	31.	AGS 111	Germplasm accession
5.	JS 20-94	JS 97-52 x JS 20-02	32.	EC457286	Germplasm accession
6.	JS 93-05	Selection from PS 73-22	33.	MACS725	JS93-05X MAUS71
7.	JS 20-116	JS 97-52 x JSM 120 A	34.	SP 37	Not known selection
8.	JS 95-60	Selection from PS 73-22	35.	NRC -125	EC54688xps1044
9.	JS 97-52	PK 327 x L 129	36.	NRC-132	JS97-52X PI086023
10.	JS 20-84	JS 98-63 x PK 768	37.	NRC-134	NRC7XAGS191
11.	JS 20-34	JS 98-63 x PK 768	38.	NRC SL-1	JS335XSL525
12.	JS 20-71	JS 97-52 x JS 90-5-12-1	39.	PS 1092	PS1042 x MACS 450
13.	RVS 2007-6	JS 20-10 x MAUS162	40.	PS 1613	PS1225XPS1042
14.	RVS 2011-35	JS 335 X PK 1042	41.	AMS 2014-1	AMS99-33XH6P5
15.	RVS 2001-4	JS 93-01x EC 390981	42.	KDS 992	JS93-05XEC241780
16.	RVS -14	JS 93-05x EC 390981	43.	VLS -94	VL Soya59X VS2005-1
17.	RVS -24	J.P 120 x JS 335	44.	SKF-SPS -11	Not known selection
18.	RVS -18	JSM110XJSM66	45.	RVS 76	MAUS-162XJSM-66
19.	NRC- 76	NRC-37XL-27	46.	NRC127	JS97-52XPI542044
20.	NRC -86	RKS15XEC481309	47.	KDS980	JS93-05XAMS1
21.	NRC- 130	EC390977XEC538828	48.	G-29	Germplasm
22.	NRC -131	EC390977XEC538828	49.	RSC-10-70	JS335X Bragg
23.	NRC -147	Germplasm accessions C210	50.	RSC-10-71	Bragg XJS335
24.	AMSMBC -18	Mutant of Bragg	51.	NRC-2	Induced mutant of Bragg
25.	AMS-100-39	Mutant of JS93-05	52.	MACS-15-20	NRC37XMohetta
26.	MACS – 1520	EC241780XMACS330	53.	MACS-58	JS2 x Improve pelican
27.	MACSNRC-1575	PI542044XJS9305			

The field experiment was conducted at the experimental field and the laboratory work at Biochemical Analysis Laboratory, Department of Plant Molecular Biology and Biotechnology, College of Agriculture, Rajmata Vijayaraje Scindia Agricultural University, Gwalior, Madhya Pradesh, India during *Kharif* 2018-19. The investigational region covered was fairly identical in terms of topography and fertility. Gwalior has subtropical, semi-arid climate and chilly winters with random showers. The average rainfall was about 312.0 mm in July, 190.6 mm in August, 166.4 mm in September and 0.0 mm in the month of October respectively. Crop was sown on July 2019. Between 60th to 70th days of crop growth season neither rain has been received nor has irrigation given manually. Leaves were collected after 70 days of sowing from five random selected plants of each line for the analysis of diverse biochemical parameters. Completely Randomized Design (CRD) was adopted and the data were analyzed as per method suggested by Snedecor and Cochran (1967).

Biochemical analysis

Photosynthetic pigments (chlorophyll a, b and total) were quantified using UV-VIS spectrophotometer at 470, 645 and 663 nm absorbance and calculated according to Arnon's equation (1949). Proline content in leaves was determined as suggested by Bates *et al.* (1973). Estimation of sugar content (mg g⁻¹ fresh weight) was estimated as methods adopted by Kachare (2017) in soybean. Malondialdehyde test (Lipid peroxidation assay, nmol g⁻¹FW) estimation was exercised with the help of method developed by Stewart and Bewley (1980). Membrane stability index (MSI) was worked out as formulae suggested by Razzaq *et al.* (2013). Total seed protein content was estimated by the process of Bradford (1976).

Determination of antioxidant enzyme activity (catalase, glutathione reductase and peroxidase)

Sample preparation was done according to the method adopted by Kachare (2017). Ascorbate peroxidase (APX) activity (EC 1.11.1.11) was computed as per method of Nakano and Asada (1981). Catalase (CAT) activity (EC 1.11.1.6) was analyzed by the UV process as proposed by Aebi (1983). Glutathione reductase (GR) activity (EC 1.6.4.2) was estimated by the technique as explained by Smith *et al.* (1988). Guaiacol peroxidase (PDX)

activity (EC 1.11.1.7) was computed by estimating the oxidation of guaiacol as protocol described by Rao *et al.* (1996).

Protein's characterization

The extracted protein was used to analyze through SDS-PAGE method according to the steps suggested by Laemmli (2011).

Results

Biochemical parameters

Variations among 53 soybean genotypes for studied biochemical traits with analysis of variance are shown in Table 2 which indicated that the alteration in their response against drought stress. Total Chlorophyll content in mgml^{-1} varied considerably in ranged of 38.13-59.87 mgml^{-1} , with maximum in genotype JS 97-52 (59.87 mgml^{-1}) chased by genotypes: RVS-14 (56.29 mgml^{-1}), JS20-94 (55.88 mgml^{-1}) and NRC-76 (55.07 mgml^{-1}). Whereas the minimum was documented in genotype JS20-84 (38.13 mgml^{-1}) pursued by genotypes RVS2011-35 (40.01 mgml^{-1}), NRC-130 (41.21 mgml^{-1}) and MACS-1520 (41.58 mgml^{-1}). Proline content varied significantly in ranged of 66.25-111.40 μgg^{-1} with highest in genotype JS 97-52 (111.40 μgg^{-1}), tracked by genotypes RVS-14 (103.45 μgg^{-1}), JS 95-60 (93.95 μgg^{-1}) and RVS 2001-4 (91.54 μgg^{-1}). However, the least showed in genotype JS335 (66.25 μgg^{-1}) followed by genotypes: JS 20-29 (67.90 μgg^{-1}), NRC-76 (68.05 μgg^{-1}), NRC-134 (68.15 μgg^{-1}) and SL-1123 (68.30 μgg^{-1}). Total Sugar in mgg^{-1} varied significantly between 2.35-5.45 mgg^{-1} with utmost in genotype JS97-52 (5.45 mgg^{-1}) which is pursued by genotypes RVS-14 (5.15 mgg^{-1}) and JS95-60 (5.15 mgg^{-1}) while lowest in genotype KDS992 (2.35 mgg^{-1}), tracked by genotype RSC-10-71 (2.45 mgg^{-1}). MDA content in nmole gm^{-1} differed significantly in ranged of 42.65 to 60.80 with greatest in genotype JS 97-52 (60.80 nmole gm^{-1}) tracked by JS95-60 (57.22 nmole gm^{-1}) and least amount (42.65 nmole gm^{-1}) in genotype RVS2007-6. Membrane stability is one of the major components in tolerance under stress. Highest membrane stability (MS) was texted in genotype JS97-52 (64.50 %) intimately trailed by genotypes RVS-14 (63.00%) and JS95-60 (61.00%), while minimum (29%) in genotype RSC-10-70. Protein synthesized in immature seeds varied significantly among 53 soybean genotypes in ranged of 34.4%-39.3% with upmost in genotype JS20-71 (39.30

%) closely pursued by genotypes: AMS-100-39 (39.10%) and MACS725 (38.6%). Whilst the lowest amount was documented in genotype JS95-60 (34.4%) tracked by genotypes JS 20-69 (34.5%) and SP 37 (34.8%).

Biochemical parameters based hierarchical cluster analysis

The hierarchical cluster analysis and the content values are presented in Table 2, and the dynamic expression profile was determined and is shown in Figure 1. Multivariate analysis based on diversity was performed using the UPGMA. The mean value of biochemical parameters of different genotypes falling in each cluster was presented in the generated dendrogram for distinguished into two major clusters *i.e.*, I and II. Cluster I divided into subclusters. These clusters further subdivided into minor clusters. Cluster I consisted genotypes RVS-2001-4 and KDS-980 as an out group and cluster II also consisted two genotypes *viz*, JS-97-52 and RVS-14 as an outgroup. The biochemical parameters illustrated similar pattern in both of these two genotypes, and they were clustered together.

Principal component analysis (PCA) of biochemical parameters

Principal component analysis (PCA) was done by considering biochemical variables simultaneously. The pattern of variations illustrated by the PCA described by correlation coefficients determined for pair-wise association of the traits. Genotypes JS-97-52 and RVS-14 situated at the unique position of the plot. The PCA correlation depicted those genotypes possessed higher and lower magnitudes of biochemical parameters occupying unique position towards the graph (Figure 2).

Activities of antioxidant enzymes

The analysis of variance (Table 2 and Figure 3) clearly indicated presence of ample variations among 53 soybean genotypes for all antioxidant enzymes activities. APX was found to be maximum in genotype JS97-52 (2.21-unit mg^{-1} protein min^{-1}) whereas minimum (0.23-unit mg^{-1} protein min^{-1}) was documented in genotype JS20-94. Catalase activity was found to be highest (0.97-unit mg^{-1} protein min^{-1}) in genotype JS 97-52 while lowest (0.30-unit mg^{-1} protein min^{-1}) was observed in genotype NRC-132. Highest Gluthione reductase (GR) activity was texted in genotype JS 97-52 (0.87-unit mg^{-1} protein

min⁻¹) narrowly pursued by genotypes: JS 95-60 (0.80 unit mg⁻¹protein min⁻¹), however, least in RVS-18 (0.22 unit mg⁻¹protein min⁻¹). Guaiacol peroxidase (PDX) ranged between 0.24-1.91 unit mg⁻¹ protein min⁻¹ with greatest (1.91 unit mg⁻¹ protein min⁻¹) in genotype JS 97-52 and lowest (0.24 unit mg⁻¹ protein min⁻¹) in genotype JS20-98. Further, elevated PDX activity was examined in genotypes *i.e.*, JS 97-52, RVS 2001-4, NRC-76 and RVS-14.

Antioxidant activities based cluster analysis

In antioxidant enzymes data based dendrogram, the genotypes were separated into two clusters *i.e.*, major and minor cluster. The major cluster contained 49 genotypes while the minor cluster had only 4 genotypes *namely*: RVS-24, JS20-84, RVS-14 and JS97-52. The major cluster was further divided into two groups. The major group is consisted 47 genotypes while minor group had only two genotypes *viz.*, NRC-76 and RVS2001-4. The major group was further divided into two sub-groups. Major sub-group contained 38 genotypes and minor sub-group consisted nine genotypes including AMS2014-1, PS-1613, VLS-94, PS-10-92, RSC-10-70, NRC-SL-1, NRC-134, NRC-132 and NRC-125. Among these nine genotypes utmost resemblance was documented between AMS2014-1 and PS-1613 and both genotypes grouped together (Figure 4).

Protein profiling through SDS-PAGE

In this experimentation, we analyzed immature seed protein pattern for fifty-three soybean genotypes using SDS-PAGE (Figure 5). Dendrogram was generated on the basis of banding pattern and studied genotypes were divided into two clusters (Figure 6). The major cluster is consisted 50 genotypes while minor cluster had only three genotypes, *viz.*, SL-1068, AGS-111 and NRC-132. Among these three genotypes, NRC-132 was found to be diversified and grouped distantly from rest of the two genotypes. The data were also analyzed on the basis of band intensity. The variations in the intensity of bands with similar molecular weight indicated the expression capability of genotypes of same protein in different genotypes.

Table 2. Mean performance of different biochemical parameters and anti-oxidant enzymatic activities of soybean genotypes

S. No.	Parameter Genotypes	Total Chlorophyll Content (mg/ml)	Proline (µg/g)	Total Sugar (mg/g)	MDA (n mole/g m)	MSI (%)	Protein (%)	Ascorbate peroxidase (unit/mg protein/min)	Catalase (unit/mg protein/min)	Glutathione reductase (unit/mg protein/min)	Guaiacol peroxidase (unit/mg protein/min)
1.	JS 20-29	53.68	67.90	3.15	46.70	50.50	36.2	0.57	0.74	0.43	0.32
2.	JS 20-69	49.90	70.35	2.70	43.57	45.00	34.5	0.40	0.72	0.30	0.51
3.	JS 335	52.30	66.25	2.75	45.39	50.5	35.6	0.58	0.40	0.29	0.62
4.	JS 20-98	52.79	71.45	3.00	47.35	37.00	37.2	0.67	0.53	0.36	0.24
5.	JS 20-94	55.88	90.40	3.90	45.17	40.50	38.1	0.23	0.44	0.41	0.42
6.	JS 93-05	50.06	75.65	3.00	44.67	36.50	37.4	0.45	0.50	0.27	0.65
7.	JS 20-116	52.39	76.05	3.75	42.73	48.00	36.1	0.31	0.64	0.27	0.75
8.	JS 95-60	47.55	93.95	5.15	57.22	61.00	34.4	0.27	0.43	0.80	0.60
9.	JS 97-52	59.87	111.4	5.45	60.80	64.50	35.7	2.21	0.97	0.87	1.91
10.	JS 20-84	38.13	81.05	4.55	46.19	49.50	36.8	1.79	0.80	0.30	0.72
11.	JS 20-34	47.13	70.50	4.30	47.07	40.50	37.1	0.26	0.85	0.34	0.39
12.	JS 20-71	50.25	77.80	3.75	46.66	39.50	39.3	0.29	0.57	0.52	0.45
13.	RVS 2007-6	47.48	70.95	2.80	42.65	53.50	37.2	0.33	0.44	0.26	0.71
14.	RVS 2011-35	40.01	81.35	3.05	45.88	43.50	38.2	0.46	0.47	0.32	0.50
15.	RVS 2001-4	41.74	91.50	2.75	47.78	45.00	36.8	0.59	0.43	0.76	1.65
16.	RVS -14	56.29	103.45	5.15	52.93	63.00	38.1	1.70	0.86	0.80	1.24
17.	RVS -24	50.39	77.65	3.00	55.12	38.50	36.2	2.04	0.64	0.27	0.37
18.	RVS -18	47.10	79.85	3.30	53.67	46.50	35.7	0.50	0.72	0.22	0.45
19.	NRC- 76	55.07	68.05	4.65	46.87	35.50	37.1	0.32	0.96	0.77	1.58
20.	NRC -86	47.20	82.00	2.65	48.20	44.50	37.8	0.47	0.63	0.63	0.57

Table 2 (Con.)

S. No.	Parameter Genotypes	Total Chlorophyll Content (mg/ml)	Proline ($\mu\text{g/g}$)	Total Sugar (mg/g)	MDA (nmole/gm)	MSI (%)	Protein (%)	Ascorbate peroxidase (unit/mg protein/min)	Catalase (unit/mg protein/min)	Glutathione reductase (unit/mg protein/min)	Guaiacol peroxidase (unit/mg protein/min)
21.	NRC-130	41.21	78.95	2.60	45.29	39.50	36.5	0.48	0.64	0.60	0.59
22.	NRC-131	52.46	87.30	3.95	43.20	51.00	37.2	0.40	0.52	0.52	0.40
23.	NRC-147	46.95	87.90	3.70	53.93	36.50	36.2	0.30	0.55	0.46	0.78
24.	AMSMBC-18	47.81	81.25	3.80	55.08	48.00	38.1	0.26	0.76	0.36	0.32
25.	AMS-100-39	49.09	80.50	3.75	46.61	44.50	39.1	0.24	0.40	0.63	0.51
26.	MACS-1520	41.58	70.95	3.15	45.76	55.00	37.4	0.55	0.61	0.58	0.79
27.	MACSNR C-1575	48.02	86.80	3.50	45.62	57.50	37.9	0.43	0.55	0.49	0.57
28.	RSC-10-52	46.07	70.15	3.05	50.20	45.00	38.1	0.41	0.35	0.40	0.40
29.	SL-1123	44.20	68.30	2.65	52.71	32.50	36.2	0.34	0.64	0.39	0.61
30.	SL-1068	43.08	91.50	4.20	54.80	53.00	35.7	0.28	0.37	0.32	0.57
31.	AGS 111	46.63	87.80	4.75	50.58	49.00	37.4	0.42	0.46	0.52	0.85
32.	EC457286	49.68	83.45	4.25	54.56	31.00	37.1	0.38	0.44	0.55	0.71
33.	MACS725	51.59	88.75	3.90	45.41	31.00	38.6	0.32	0.50	0.45	0.90
34.	SP 37	48.51	81.30	3.55	46.09	41.00	34.8	0.43	0.69	0.40	0.53
35.	NRC-125	51.42	85.60	4.05	50.55	50.50	37.6	0.87	0.43	0.46	0.63
36.	NRC-132	47.08	85.75	3.25	52.92	57.00	37.9	0.77	0.30	0.62	0.60
37.	NRC-134	52.14	68.15	3.70	45.33	32.50	37.1	0.82	0.36	0.60	0.64
38.	NRC SL-1	41.69	82.30	2.90	51.42	40.50	38.1	0.80	0.63	0.60	0.63
39.	PS 1092	46.87	90.85	2.60	46.63	52.00	37.2	0.82	0.39	0.24	0.81
40.	PS 1613	49.63	78.30	3.00	43.53	51.00	38.4	0.70	0.44	0.49	0.87

Table 2 (Con.)

S. No.	Parameter Genotypes	Total Chlorophyll Content (mg/ml)	Proline (µg/g)	Total Sugar (mg/g)	MDA (n mole/g m)	MSI (%)	Protein (%)	Ascorbate peroxidase (unit/mg protein/min)	Catalase (unit/mg protein/min)	Glutathione reductase (unit/mg protein/min)	Guaiacol peroxidase (unit/mg protein/min)
41.	AMS 2014-1	45.19	80.00	2.75	45.60	51.50	37.2	0.76	0.43	0.47	0.92
42.	KDS 992	50.18	70.35	2.35	45.13	34.50	36.5	0.44	0.59	0.34	0.85
43.	VLS -94	48.38	82.00	3.05	50.68	37.00	36.2	0.63	0.39	0.25	0.80
44.	SKF-SPS -11	40.42	85.30	3.35	53.92	51.00	37.4	0.51	0.43	0.31	0.73
45.	RVS 76	42.42	71.65	2.95	54.92	51.00	36.8	0.68	0.52	0.36	0.58
46.	NRC127	48.76	70.65	2.50	54.91	52.00	37.1	0.44	0.50	0.39	0.46
47.	KDS980	42.19	71.45	2.95	55.23	41.50	35.7	0.47	0.57	0.48	0.72
48.	G-29	44.14	78.55	2.60	55.10	47.50	36.2	0.40	0.45	0.36	0.55
49.	RSC-10-70	46.64	76.00	3.00	54.78	29.00	37.1	0.80	0.64	0.54	0.56
50.	RSC-10-71	51.00	71.55	2.45	44.11	50.50	36.4	0.52	0.60	0.59	0.73
51.	NRC-2	45.78	75.55	2.80	56.41	50.50	37.2	0.67	0.42	0.32	0.60
52.	MACS-15-20	50.98	77.70	2.70	56.37	58.50	37.9	0.62	0.49	0.36	0.47
53.	MACS-58	51.90	81.05	2.85	45.21	48.00	36.5	0.45	0.33	0.45	0.33
	SE (m)	1.076566	1.130809	0.154004	3.160806	0.845032	0.147046	0.009329	0.017895	0.014916	0.016474
	CD_{0.05}	3.057168	3.211204	0.437332	8.97587	2.399671	0.417574	0.026	0.050817	0.042358	0.046783
	Range	38.13-59.87	66.25-111.40	2.35-5.45	42.65-60.80	29.00-64.50	34.4-39.3	0.23 - 2.21	0.30-0.97	0.22-0.87	0.24-1.91




Figure 1. Biochemical based hierarchical cluster analysis of soybean genotypes based on different biochemical parameters (total chlorophyll content, proline, total sugar, MDA, MSI and protein %)




Figure 3. Anti-oxidative enzymes activities of different genotypes




Figure 4. Dendrogram showing relationship among soybean genotypes based on different antioxidative enzymes (Ascorbate peroxidase, Catalase, Glutathione reductase and Guaiacol peroxidase)




Figure 5. SDS protein profiling of 53 soybean genotypes




Figure 6. Dendrogram showing relationship among different soybean genotypes based on SDS- Profiling

Discussion

Biochemical parameters

Generally, the level of chlorophyll content in leaves determines the rate of photosynthesis. In the current study, a reduction in chlorophyll component was found in susceptible genotypes in comparison to tolerant genotypes. Similar results in soybean were found earlier by Hossain *et al.* (2015). Reduced level of chlorophyll synthesis in susceptible genotypes may be the reason of less activity of the photosynthetic elements. Previously, loss of chloroplast membranes under drought stress has also been documented by Anjum *et al.* (2011). Parallel reduction in chlorophyll levels in several other plant species *viz.*, soybean, maize, rice, chickpea, pearl millet *etc.* have been reported (Zhang *et al.*, 2007; Sahu *et al.*, 2020; Choudhary *et al.*, 2021; Sharma *et al.*, 2021). High reduction in chlorophyll content was found in drought susceptible genotypes in the present study.

Role of proline in osmotic regulation under water stress has been monitored in various plant species (Rengasamy, 2002; Guo *et al.*, 2009). Soybean genotypes with significant rise in proline contents have been considered as drought tolerant. Highest proline content increment was found in genotype JS97-52 followed by genotypes RVS2001-4 and JS95-60. Previously, Khan *et al.* (2015), Kachare *et al.* (2019) and Sharma *et al.* (2021) observed comparable trend in the proline content during their study on screening of soybean genotypes tolerant to drought stress. Increased proline content maintains cell water level under drought (Ghorbanli *et al.*, 2012; Choudhary *et al.*, 2021). Further, George *et al.* (2015) suggested that increased proline has osmoprotective functions by preventing separation of enzymes during metabolic activities.

Among osmotic regulating substances, total soluble sugars are noteworthy contributors of drought tolerance (Gurrieri *et al.*, 2020; Choudhary *et al.*, 2021). During present investigation, genotype JS97-52 had maximum soluble sugar tracked by genotypes RVS2001-4 and JS95-60. These findings indicate the presence of possible drought tolerance in these genotypes. Previously, Kachare *et al.* (2019) and Sharma *et al.* (2021) also observed momentous enhancement in total soluble sugar content under drought stress while studying on Indian soybean genotypes.

Lipid peroxidation mechanism is a cursor of oxidative stress under water stress. Malondialdehyde (MDA) content is considered as a sign of membrane lipid peroxidation and shows the level of damage in membrane underneath stress (Wang *et al.*, 2019). MDA content in nmole gm⁻¹ was found

greatest in genotype JS 97-52. Genotypes exhibited higher MDA content in leaves might be tolerant against drought (Tatar and Gevrek, 2008) due to more synthesis of Reactive Oxygen Species in comparison of the rest of the genotypes and this parameter can be applicable to find out drought tolerance in genotypes (Chug *et al.*, 2011).

According to Blackman *et al.* (1995) the increments in MSI indicates the reduction of lipid peroxidation with oxidative bursts under water stress conditions. In previous studies, MSI has been used frequently for screening of drought tolerance in crop species (Farooq and Azam, 2002). In similar studies, Almeselmani *et al.* (2012), Kachare *et al.* (2019) and Sharma *et al.* (2021) reported significant genotypic differences in membrane stability as an indirect criterion for selection of drought tolerance in soybean.

Anti-oxidant enzymes

Anti-oxidative enzymes have vital role in the protection of plants in water stress due to their contribution in tolerance mechanism (Blokhina *et al.*, 2003; Xue *et al.*, 2011). This may be entire capacity of a particular genotype to combat against these stresses. Previous reports proved the role of antioxidant enzymes in the mechanism of drought and dehydration tolerance in soybean (Vasconcelos *et al.*, 2009; Xue *et al.*, 2011). Major system in plants under abiotic stress for detoxification of hydrogen peroxide is the ascorbate-glutathione cycle in which Ascorbate peroxidase (APX) helps in the conversion of H₂O₂ into H₂O, in chloroplast (Mittler and Zilinskas, 1994; Correa-Aragunde *et al.*, 2013). Enhanced APX activity was observed in genotypes JS97-52 followed by RV-24 as compared to rest of the genotypes possibly due to appearance of prominent H₂O₂ detoxification with prevention of H₂O₂-mediated cell damage (Kommavarapu *et al.*, 2013). Increased APX activity in soybean under drought stress has been earlier reported by Kausar *et al.* (2012), Kachare *et al.* (2019) and Sharma *et al.* (2021). Catalase (CAT) enzymes are capable to alter millions of H₂O₂ molecules into H₂O and O₂ in a second (Chelikani *et al.*, 2004). Enhanced CAT and GR activity was found in soybean genotype JS97-52. In an earlier research, conducted by Masoumi *et al.* (2011) a positive correlation between drought tolerance in plant genotypes and enhanced CAT was reported. Similarly, Liu *et al.* (2013), Kachare *et al.* (2019) and Sharma *et al.* (2021). also observed a significant increase in GR activity in some soybean genotypes. Earlier, Porcel *et al.* (2003) reported increased GR activity in drought tolerant soybean genotypes. This indicates the enhanced ROS scavenging ability of drought tolerant plants as a result negligible injury to plants under stress. Role of GR as an important enzyme to sustain redox

condition of plant by converting oxidized glutathione (GSSG) into reduced glutathione (GSH) with the help of NADPH has also been proved by Garg *et al.* (2012). Enhanced PDX activity was evidenced in genotype JS97-52 while lowest was in genotype JS20-98. This indicates the possible tolerance in these genotypes as earlier stated by Murthy *et al.* (2012). Similar to the present research, Kachare (2017) and Sharma *et al.* (2021) also noticed directly proportional relation between PDX activity and level of water stress in soybean. Akitha-Devi *et al.* (2015) reported enhanced activities of antioxidant enzymes with reduction in osmotic potential. These findings clarify the relation between activity of antioxidant enzyme and defense mechanisms against water stress in soybean.

Protein profiling

Soybean seed contains about 35-40% proteins with big contribution of essential amino acids. It is also a rich source of antioxidants and unsaturated fatty acids. Many of these proteins have functions in the management of water deficit (Qayyum *et al.*, 2011). These proteins play major role in uptake of water from environment for the betterment of plant health. So, the proteins help the plants in completion of growth and development cycle normally. On average, 12 bands per genotype were detected in a range of molecular weight from 3.5 to 43.0 kDa in present study correspondingly to Arumingtyas and Savitri (2014) and Kachare (2017). Sahu *et al.* (2020) and Gupta *et al.* (2021) also detected range of protein bands in chickpea.

It is concluded that the basis of different biochemical parameters and anti-oxidant enzymes activities genotypes *viz.*, JS97-52, RVS-14 and JS95-60 were found with drought tolerance. The findings of the present study provide a base for the use of these soybean genotypes for further hybridization to develop drought tolerant varieties. These results also open the door for the applications of advanced biotechnological tools for deep analysis of drought tolerance mechanism in soybean crop.

References

- Aebi, H. E. (1983). Catalase. In "method of enzymatic analysis", VCH, Weinheim, Germany-Deerfield, FL, 3:273-286.
- Akitha-Devi, M. K. and Giridhar, P. (2015). Variations in physiological response, lipid peroxidation, antioxidant enzyme activities, proline and isoflavones content in soybean varieties subjected to drought stress. Proceedings of the National Academy of Sciences, 85:35-44.

- Almeselmani, M., Saud, A., Al-Zubi, K., Abdullah, F., Hareri, F., Nassan, M., Ammar, M. A. and Kanbar, O. (2012). Physiological performance of different durum wheat varieties grown under rainfed condition. *Global Journal of Science Frontier Research*, 12:55-63.
- Anjum, S. A., Xie, X., Wang, L., Saleem, M. F., Chen, M. and Wang, L. (2011). Morphological, physiological and biochemical responses of plants to drought stress. *African Journal of Agricultural Research*, 6:2026-2032.
- Arnon, D. I. (1949). Copper enzymes in isolated chloroplasts, polyphenoxidase in *Betavulgaris*. *Plant Physiology*, 24:1-15.
- Arumingtyas, E. L. and Savitri, E. S. (2014). Identification and characterization of drought stress protein on soybean (*Glycine max* L. Merr). *Research Journal of Pharmaceutical, Biological and Chemical Sciences*, 5:789-796.
- Bates, L. S., Waldren, R. P. and Teare, I. D. (1973). Rapid determination of free proline for water-stress studies. *Plant and Soil*, 39:205-207.
- Blackman, S. A., Obendorf, R. L. and Leopold, A. C. (1995). Desiccation tolerance in developing soybean seeds: The role of stress proteins. *Plant Physiology*, 93:630-638.
- Blokhina, O., Virolainen, E. and Fagerstedt, K. V. (2003). Antioxidants, oxidative damage and oxygen deprivation stress: a review. *Annals of Botany*, 91:179-194.
- Bradford, M. M. (1976). A rapid and sensitive method for the quantitation of microgram quantities of protein utilizing the principle of protein-dye binding. *Analytical Biochemistry*, 72:248-254.
- Buezo, S. Á., Moran, J. F., Soba, D. and Aranjuelo, I. (2018). Drought tolerance response of high-yielding soybean varieties to mild drought: physiological and photochemical adjustments. *Physiologia Plantarum*, <https://doi.org/10.1111/ppl.12864>.
- Chelikani, P., Fita, I. and Loewen, P. C. (2004). Diversity of structures and properties among catalases. *Cellular and Molecular Life Sciences*, 61:192-208.
- Choudhary, M. L., Tripathi, M. K., Tiwari, S., Pandya, R. K., Gupta, N., Tripathi, N. and Parihar, P. (2021). Screening of pearl millet [*Pennisetum glaucum* (L.) R. Br.] germplasm lines for drought tolerance based on morpho-physiological traits and SSR markers. *Current Journal of Applied Science and Technology*, 40:46-63.
- Correa-Aragunde, N., Foresi, N., Delledonne, M. and Lamattina, L. (2013). Auxin induces redox regulation of ascorbate peroxidase 1 activity by S-nitrosylation/denitrosylation balance resulting in changes of root growth pattern in *Arabidopsis*. *Journal of Experimental Botany*, 64:3339-49.
- Chug, V., Kaur, N. and Gupta, A. K. (2011). Role of antioxidant and anaerobic metabolism enzymes in providing tolerance to maize (*Zea mays* L.) seedlings against water logging. *Indian Journal of Biochemistry & Biophysics*, 48:346-52.
- Dhanda, S. S., Sethi, G. S. and Behl, R. K. (2004). Indices of drought tolerance in wheat genotypes at early stages of plant growth. *Journal of Agronomy and Crop Science*, 190:6-12.
- Farooq, S. and Azam, F. (2002). Co-existence of salt and drought tolerance in Triticaceae. *Hereditas*, 135:205-210.
- Gardner-Outlaw, T. and Engelman, R. (1997). Sustaining water, easing scarcity: A second update. Washington, USA: Population Action International.
- Garg, B., Jaiswal, J. P., Misra, S., Tripathi, B. N. and Prasad, M. (2012). A comprehensive study on dehydration-induced antioxidative responses during germination of Indian bread wheat (*Triticum aestivum* L. em Thell) cultivars collected from different agroclimatic zones. *Physiology and Molecular Biology of Plants*, 18:217-228.

- George, S., Minhas, N. M., Jatoi, S. A., Siddiqui, S. U. and Ghafoor, A. (2015). Impact of polyethylene glycol on proline and membrane stability index for water stress regime in tomato (*solanum lycopersicum*). Pakistan Journal of Botany, 47:835- 844.
- Ghorbanli, M., Gafarabad, M., Amirkian, T. and Mamaghani, B. A. (2012). Investigation of praline, total protein, chlorophyll, ascorbate and dehydro-ascorbate changes under drought stress in Akria and Mobil tomato cultivars. Iranian Journal of Plant Physiology, 3:651-658.
- Gupta, N., Tiwari, S. Tripathi, M. K. and Bhagyawant, S. S. (2021). Antinutritional and protein-based profiling of diverse desi and wild chickpea accessions. Current Journal of Applied Science and Technology, 40:7-18.
- Gurrieri, L., Merico, M., Trost, P., Forlani, G. and Sparla, F. (2020). Impact of drought on soluble sugars and free proline content in selected Arabidopsis mutants. Biology, 9:367; doi:10.3390/biology9110367.
- Guo, P., Baum, M., Grando, S., Ceccarelli, S., Bai, G., Li, R., Korff, M., Varshney, R. K., Graner, A. and Valkoun, J. (2009): Differentially expressed genes between drought-tolerant and drought-sensitive barley genotypes in response to drought stress during the reproductive stage. Journal of Experimental Botany, 60:3531-3544.
- Hossain, M. M., Lam, H. M. and Zhang, J. (2015). Responses in gas exchange and water status between drought-tolerant and susceptible soybean genotypes with ABA application. The Crop Journal, 3:500-506.
- Kachare, S. (2017). Studies on morpho-physiological changes and gene expression under drought condition in soybean [*Glycine max* (L.) Merrill]. (Ph. D Thesis), JNKVV, Jabalpur.
- Kachare, S., Tiwari, S., Tripathi N. and Thakur V. V. (2019). Assessment of genetic diversity of soybean (*Glycine max* (L.) Merr.) genotypes using qualitative traits and microsatellite makers. Agricultural Research, DOI: 10.1007/s40003-019-00412-y.
- Kausar, R., Hossain, Z., Makino, T. and Komatsu, S. (2012). Characterization of ascorbate peroxidase in soybean under flooding and drought stresses. Molecular Biology Reports, 39:10573-10579.
- Kommavarapu, M., Balaraju, P., Ramakrishna, B. and Ram Rao, S. S. (2013). Effect of brassinosteroids on germination and seedling growth of radish (*Raphanus sativus* L.) under PEG-6000 induced water Stress. American Journal of Plant Sciences, 4:2305-2313.
- Khan, S. A., Karim, M. A., Mahmud, Abullah-Al, Sarveen, S., Bazzaz, M. M. and Hossain, M. A. (2015). Plant water relations and proline accumulations in soybean under salt and water stress environment. Journal of Plant Sciences, 3:272-278.
- Laemmli (2011). Laemmli-SDS-PAGE. Bio Protocol, 1(11) DOI:10.21769/BioProtoc.80.
- Liu, J., Wang, X., Hu, Y., Hu, W. and Yurong, B. (2013). Glucose-6-phosphate dehydrogenase plays a pivotal role in tolerance to drought stress in soybean roots. Plant Cell Reports, 32:415-429.
- Masoumi, H., Farrokh, D., Jahanfar, D., Ghorban, N. and Davood, H. (2011). Chemical and biochemical responses of soybean (*Glycine max* L.) cultivars to water deficit stress. Australian Journal of Crop Science, 5:544-553.
- Mittler, R. and Zilinskas, B. A. (1994). Regulation of pea cytosolic ascorbate peroxidase and other antioxidant enzymes during the progression of drought stress and following recovery from drought. Plant Journal, 5:397-405.
- Mishra, N., Tripathi, M. K., Tiwari, S., Tripathi, N. and Trivedi, H. K. (2020). Morphological and molecular screening of soybean genotypes against yellow mosaic virus disease. Legume Research an International Journal, DOI: 10.18805/LR4240.

- Mishra, N., Tripathi, M. K., Tiwari, S., Tripathi, N., Ahuja, A., Sapre, S. and Tiwari, S. (2021a). Cell suspension culture and *in vitro* screening for drought tolerance in soybean using poly-ethylene glycol. *Plants*, 10:517-536.
- Mishra, N., Tripathi, M. K., Tiwari, S., Tripathi, N., Gupta, N. and Sharma, A. (2021b). Morphological and physiological performance of Indian soybean [*Glycine max* (L.) Merrill] genotypes in respect to drought. *Legume Research an International Journal*, DOI:10.18805/LR-4550.
- Murthy, S. M., Devaraj, V. R., Anitha, P. and Tejavathi, D. H. (2012). Studies on the activities of antioxidant enzymes under induced drought stress in *in vivo* and *in vitro* plants of *Macrotyloma uniflorum* (Lam.) Verdc. *Recent Research in Science and Technology*, 4:34-37.
- Nakano, Y. and Asada, K. (1981). Hydrogen peroxide is scavenged by ascorbate-specific peroxidase in spinach chloroplasts. *Plant Cell Physiology*, 22:867-880.
- Porcel, R., Barea, J. M. and Ruiz-Lozano, J. M. (2003). Antioxidant activities in mycorrhizal soybean plants under drought stress and their possible relationship to the process of nodule senescence. *New Phytologist*, 157:135-143.
- Qayyum, A., Razzaq, A., Ahmad, M. and Jenks, M. A. (2011). Water stress causes differential effects on germination indices, total soluble sugar and proline content in wheat (*Triticum aestivum* L.) genotypes. *African Journal of Biotechnology*, 10:64. Doi.14038-14045.
- Rao, M., Paliyath, G. and Ormrod, D. P. (1996). Ultraviolet-B and ozone induced biochemical changes in antioxidant enzymes of *Arabidopsis thaliana*. *Plant Physiology*, 110:125-136.
- Razzaq, A., All, Q., Qayyum, A., Mahmood, I., Ahmad, M. and Rasheed, M. (2013). Physiological responses and drought resistance index of nine wheat (*Triticum aestivum* L.) cultivars under different moisture conditions. *Pakistan Journal of Botany*, 45:151-155.
- Rengasamy, P. (2002). Transient salinity and subsoil constraints to dryland farming in Australian sodic soils: an overview. *Australian Journal of Experimental Agriculture*, 42:351-361.
- Sahu, V. K., Tiwari, S., Gupta, N., Tripathi, M. K. and Yasin, M. (2020). Evaluation of physiological and biochemical contents in desi and kabuli chickpea. *Legume Research an International Journal*, 10. doi. 18805/LR-4265.
- Sharma, A., Tripathi, M. K., Tiwari, S., Gupta, N., Tripathi, N. and Mishra, N., (2021). Evaluation of soybean (*Glycine max* L.) genotypes on the basis of biochemical contents and anti-oxidant enzyme activities. *Legume Research an International Journal*, In Press.
- Smith, I. K., Vierheller, T. V. and Thorne, C. A. (1988). Assay of glutathione reductase in crude tissue homogenates using 5, 5'-dithiobis (2-nitrobenzoic acid). *Annals of Biochemistry*, 175:408-413.
- Snedecor, G. W. and Cochran, W. G. (1889). *Statistical methods*. VIII Ed. Wiley-Blackwell.
- Specht, J. E., Chase, K., Macrander, M., Graef, G. L., Chung, J., Markwell, J. P., Germann, M., Orf, J. H. and Lark, K. G. (2001). Soybean response to water. A QTL analysis of drought tolerance. *Crop Science*, 41:493-509.
- Stewart, R. R. C. and Bewley, J. D. (1980). Lipid peroxidation associated aging of soybean axes. *Plant Physiology*, 65:245-248.
- Tatar, O. and Geverek, M. N. (2008). Influence of water stress on proline accumulation, lipid peroxidation and water content of wheat. *Asian Journal of Plant Sciences*, 7:409-412. DOI:10.3923/ajps.2008. .
- Tiwari, S. and Tripathi, M. K. (2005). Comparison of morphogenic ability of callus types induced from different explants of soybean (*Glycine max* L. Merrill). *Legume Research*

- an International Journal, 28:115-118.
- Tripathi, M. and Tiwari, S. (2004). Morphogenesis and plantlet regeneration from soybean (*Glycine max* L. Merrill) leaf discs influenced by genotypes and plant growth regulators. Legume Research an International Journal, 27:88-93.
- Upadhyay, S., Singh, A. K., Tripathi, M. K., Tiwari, S. and Tripathi, N. (2020a). Validation of simple sequence repeats markers for charcoal rot and *Rhizoctonia* root rot resistance in soybean genotypes. International Journal of Applied Business Research, 10:137-144.
- Upadhyay, S., Singh, A. K., Tripathi, M. K., Tiwari, S., Tripathi, N. and Patel, R. P. (2020b). *In vitro* selection for resistance against charcoal rot disease of soybean [*Glycine max* (L.) Merrill] caused by *Macrophomina phaseolina* (Tassi) Goid. Legume Research an International Journal, DOI: 10.18805/LR-4440.
- Vasconcelos, A. C. F., Zhang, X. Z., Ervin, E. H. and Kiehl, J. D. (2009). Enzymatic antioxidant responses to biostimulants in maize and soybean subjected to drought. Scientia Agricola, 66:395-402.
- Wang, N., Yuan, M., Chen, H., Li, Z. and Zhang, M., (2019). Effects of drought stress and rewatering on growth and physiological characteristics of invasive *Aegilops tauschii* seedlings. Acta Prataculturae Sinica, 28:70-78.
- Wu, Z. and Zhang, Y. (2019). Effects of exogenous auxin on physiological and biochemical characteristics of soybean under PEG simulated drought stress. Hubei Agricultural Sciences, 58:16-19.
- Xue, G. P., Way, H. M., Richardson, T., Drenth, J., Joyce, P. A. and McIntyre, C. L. (2011). Overexpression of *TaNAC69* leads to enhanced transcript levels of stress up-regulated genes and dehydration tolerance in bread wheat. Molecular Plant, 4:697-712.
- Zhang, M., Duan, L., Tian, X., He, Z., Li, J., Wang, B. and Li, Z. (2007). Uniconazole-induced tolerance of soybean to water deficit stress in relation to changes in photosynthesis, hormones and antioxidant system. Journal of Plant Physiology, 164:709-717.

(Received: 4 May 2021, accepted: 30 June 2021)